

## LAPAROSCOPIC VS SURGICAL APPROACH OF TREATMENT FOR MECHANICAL JAUNDICE WITH PANCREATICOBILIARY MALIGNANCY

*Nurboboyev Adhambek Uygunovich*  
*Assistants Bukhara State Medical Institute*

---

### Abstract

The creation of qualitatively new laparoscopic equipment, the improvement of laparoscopic instruments required the revision of outdated provisions and the development of modern principles of laparoscopic surgery. In 2018 – 2020, 20493 operations were performed in the Bukhara branch, of which 9461 (12.1%) were cavity operations. The examined group consisted of 2,479 (26.2%) patients with urgent diseases of the abdominal cavity, among all the abdominal operations performed in the period from 2018 to 2020 using laparoscopic technologies. A palliative treatment of biliary obstruction is advisable to mitigate the effects of jaundice, independently of whether a surgical resection for attempting a cure is feasible or not. Endoscopic stent drainage has been proposed as an alternative to biliary-enteric bypass surgery to palliate malignant biliary obstruction.

**Keywords:** pancreatic biliary malignancy, laparoscopic equipment, biliary obstruction, abdominal surgery, abdominal trauma, fibro esophagogastroduodenoscopy.

---

**Relevance:** Despite the wide arsenal of non-invasive research methods, the diagnosis of acute surgical diseases of the abdominal cavity in some cases is very difficult. In this situation, the use of laparoscopy allows timely diagnosis and determination of surgical tactics. Biliary obstruction is a frequent presenting feature of pancreaticobiliary malignancy. Unfortunately, when a diagnosis of pancreaticobiliary malignancy is made the patient is usually in a terminal stage<sup>1,2</sup>. However, a palliative treatment of biliary obstruction is advisable to mitigate the effects of jaundice, independently of whether a surgical resection for attempting a cure is feasible or not. Endoscopic stent drainage has been proposed as an alternative to biliary-enteric bypass surgery to palliate malignant biliary obstruction. In addition, alternative approaches to biliary stent placement have prompted a particular interest towards a comparative evaluation aimed at assessing optimal stent material and design, as well as stent placement strategies. A comprehensive comprehensive analysis of the use of laparoscopic operations in emergency abdominal surgery and their improvement is an urgent task of modern clinical surgery. With the development of laparoscopic surgery and the accumulation of experience, the interest of many surgeons in the use of this method in emergency surgery is expanding. Within the framework of general scientific and technological progress, the constant introduction of new technologies makes them an integral part of a particular field of medicine. Today, emergency surgery is impossible to imagine without laparoscopy. In many complex cases, laparoscopy remains indispensable, as it allows you to give a direct visual assessment of the pathological process, identify its prevalence, perform a targeted biopsy and perform instrumental palpation. It is safe to say that the number of urgent operations performed by laparoscopic method is steadily increasing. This circumstance requires the development of clear recommendations on the use of laparoscopic technologies in emergency surgery. The method makes it possible to carry out diagnostics with a greater degree of informativeness, and when establishing pathology in the abdominal cavity, it provides an opportunity to transform diagnostic manipulation into therapeutic. In the Clinic of Surgical Diseases and Resuscitation of the Bukhara State Medical Institute, from 2018 to 2020, out of

9461 emergency abdominal operations performed, laparoscopic technique was used in 2479 (26.2%) observations.

The following laparoscopic operations were performed: laparoscopic cholecystectomy (LCE) for acute calculous cholecystitis was performed in 1984 (80%) patients, diagnostic and therapeutic laparoscopy - 49 (1.9%), diagnostic and therapeutic thoracoscopy in 32 (1.3%) patients, laparoscopic appendectomy - 18 (0.7%), suturing of perforative ulcers of the stomach and duodenum (duodenum) -15(0.6%)patients, laparoscopic cystectomy (gynecology)- 112(4.5%), laparoscopic tubectomy -181(7.3%), laparoscopic dissection of adhesions - 24 (0.9%) and laparoscopic ureterolithotomy - 64(2.6%) patients. Biliary obstruction results in a variety of biochemical and physiologic disturbances, such as elevated serum bilirubin, altered liver function tests, as well as impaired hepatic and renal functions with associated coagulation problems. In view of the fact that most patients with pancreatic cancer are diagnosed at an advanced and inoperable stage of the disease, the intent of treatment is palliation. Among these patients, some will survive only a few weeks, many of them will die within six months, but some others may survive in a fairly good health condition for longer time. Until recently, two kinds of therapeutic options were offered to such patients: (a) surgical bypass and (b) endoprosthesis. The percutaneous insertion of an endoprosthesis has been shown to provoke an excess morbidity<sup>19</sup> due to a 12 French gauge access through the liver, and it should not be considered as the primary option for common bile duct obstruction, although it can be recommended for the palliative treatment of porta hepatis strictures.

**The purpose of the study:** Improving the results of surgical treatment of patients with acute surgical diseases of the abdominal cavity, by improving diagnostic and therapeutic laparoscopy.

**Materials and methods of research:** The work was carried out in the Bukhara branch of the Emergency Hospital on the basis of surgical departments. In 2018 – 2020, 20493 operations were performed in the Bukhara branch, of which 9461 (12.1%) were cavity operations. The examined group consisted of 2,479 (26.2%) patients with urgent diseases of the abdominal cavity, among all the abdominal operations performed in the period from 2018 to 2020 using laparoscopic technologies.

The paper analyzes the results of a parallel comparative study of laparoscopic and open methods of surgical treatment. All the admitted patients were subjected to a comprehensive clinical study according to the generally accepted scheme using traditional laboratory and radiological methods, general clinical and biochemical studies were conducted; in-depth study of the coagulation system; overview radiography of the abdominal cavity, radiopaque studies of the gastrointestinal tract (gastrointestinal tract); computed tomography (CT), including with contrast, ultrasound examination (ultrasound) of the abdominal cavity and pelvis, magnetic resonance imaging in rare cases - to clarify the diagnosis; endoscopic studies fibroesophagogastroduodenoscopy. According to the indications, patients with cholelithiasis underwent endoscopic retrograde pancreato-cholangiography, supplemented, if necessary, by endoscopic papillosphincterotomy. The patients subjected to laparoscopic examination and surgical treatment were aged from 23 to 77 years, while there were 961 patients older than 60 years (38.7%). The following laparoscopic operations were performed: laparoscopic cholecystectomy (LCE) for acute calculous cholecystitis was performed in 1984 (80%) patients, diagnostic and therapeutic laparoscopy - 49 (1.9%), diagnostic and therapeutic thoracoscopy in 32 (1.3%) patients, laparoscopic appendectomy - 18 (0.7%), suturing of perforative ulcers of the stomach and duodenum (DPC) - 15 (0.6%) patients, laparoscopic cystectomy (gynecology) - 112 (4.5%), laparoscopic tubectomy -181 (7.3%), laparoscopic dissection of adhesions - 24 (0.9%) and laparoscopic ureterolithotomy - 64 (2.6%) patients. Table No. 1

Types of laparoscopic operations in the Bukhara branch of the Emergency Hospital 2018-2020yy

| Operation by years                      | 2018 | 2019 | 2020 | 2018-2020 |
|---|------|------|------|-----------|
| Diactostic and treatment of laparoscopy | 12   | 21   | 16   | 49(1,9%)  |
| Appendectomia via the Laporoscopy       | 5    | 6    | 7    | 18(0,7%)  |
| Cholisisectomia via the Laporoscopy     | 603  | 708  | 673  | 1984(80%) |
| Laporoscopy inthePerphroratsio ulcer    | 4    | 5    | 6    | 15(0,6%)  |
| Laparoscopic dissection of adhesions    | 4    | 8    | 12   | 24(0,9%)  |

|   |     |     |     |           |
|---|-----|-----|-----|-----------|
| Laparoscopic cystectomy (gynecology)    | 35  | 30  | 47  | 112(4,5%) |
| Laparoscopic tubectomy                  | 72  | 74  | 35  | 181(7,3%) |
| Diagnostic and therapeutic thoracoscopy | 2   | 8   | 22  | 32(1,3%)  |
| Ureterolithotomy                        |     |     | 64  | 64(2,6%)  |
| Total laparoscopic operations           | 737 | 860 | 882 | 2479      |

To assess the effectiveness of the technique of surgical intervention, the following indicators were evaluated: the duration of surgery, normalization of body temperature, the need for drug stimulation of the intestine, the recovery time of intestinal peristalsis, the time of activation of the patient, the volume of intraoperative blood loss, the duration of hospitalization, the need and duration of stay in the intensive care unit (ICU), the need and duration of the use of analgesics, the number and the severity of the intra- and postoperative complications, cosmetic defect after surgery, the need for repeated surgical interventions, the number of conversions.

Laparoscopic intervention was performed in a limited hermetic space (in the abdominal, thoracic cavity), which was created by the introduction of gas. A certain level of intra-abdominal pressure (12-15 mmHg) was created by introducing carbon dioxide into the abdominal cavity and was stably maintained throughout the operation, which provided favorable conditions for performing the necessary manipulations under the control of a laparoscope. The gas was injected through a Veresh needle by puncture of the anterior abdominal wall. The air cushion obtained in the abdominal cavity prevented damage to its organs during the introduction of trocars. A Veresh needle was inserted into the abdominal cavity through a small incision (up to 1 cm) in the umbilical region along the white line of the abdomen. To prevent damage to internal organs or vessels with a Veresh needle, the anterior abdominal wall was gripped with a brush into a longitudinal fold or a clamp and lifted up. After the creation of the pneumoperitoneum, trocars were inserted into the abdominal cavity, starting with the paraumbilical, through which a laparoscope was inserted into the abdominal cavity. Other trocars were administered under visual laparoscopic control. Laparoscopic instruments were introduced through trocars, with the help of which laparoscopic intervention was performed.

**Results and discussions:** This study was programmed in order to provide a basis for advising patients on the possibility of having their jaundice relieved either by an endoscopic stenting procedure or by a surgical bypass procedure. The prerequisites for a successful endoscopic stenting are (a) accuracy of diagnosis and (b) exclusion of patients presenting with tumors potentially treatable by a curative resection. This evaluation is simple whenever metastases are evident or whenever serious medical disorders are coincidental, thus precluding resection, but it may be difficult when a solitary primary lesion is detected.

As a result of 1984 laparoscopic cholecystectomies (LCE), complications of LCE for acute cholecystitis were observed in 14 (0.8%) patients. Parenchymal bleeding from the gallbladder bed was observed in 11 (0.6%) patients; in all, bleeding was stopped by laparoscopic coagulation. Damage to the extrahepatic bile ducts occurred in 3 (0.2%) patients and required laparotomy. In 2 observations, a lateral lesion of the choledochus was sutured. Completely crossed hepaticocholedoch 1 patients were restored on the drainage of the Kerus.

Damage to the bile ducts is facilitated by various anatomical variants of the structure and mutual disposition of the tubular structures of the hepatoduodenal ligament. An extremely dangerous situation is when the common hepatic duct is attached directly to the neck of the gallbladder, and the latter has a short duct. We have repeatedly encountered similar cases, but in one of them we could not avoid a complete intersection of the common bile duct.

Damage to the hepaticocholedocha can also occur in the absence of anatomical prerequisites, if the technique of performing the operation is violated. These are the so-called "classic" cases of duct injury. They occur when excessive traction of the gallbladder is created. In order to prevent complications in the isolation of the cystic duct and artery, we use the following technique: using an electric hook, we dissect the peritoneum from the lateral side of the gallbladder to the level of the confluence of the cystic

duct into the common hepatic duct, then we isolate the medial wall of the gallbladder with mandatory tissue preparation behind the cystic duct. Thus, the shape of the neck is created according to the type of "elephant trunk", which makes it easy to differentiate the cystic duct to the confluence with the common hepatic duct. If damage to the hepaticocholedochus is detected during surgery, it is advisable to apply its primary suture with a duct width of at least 5 mm with a sufficient wall thickness. Reconstructive surgery requires an experienced surgeon, the necessary tools and suture material.

In the postoperative period, complications occurred in 22 (1.1%) patients: bile drainage from the abdominal cavity - in 8 (0.4%), subhepatic abscess in the bed of the gallbladder — in 2 (0.1%), suppuration of epigastric puncture was observed — in 12 (0.6%) patients. (Table 2.)

**Table 2. Complications after laparoscopic cholecystectomy**

| Types of complications                           | Number of abs patients (%) |
|--|----------------------------|
| Bile drainage from the abdominal cavity          | 8 (0,4%)                   |
| Subhepatic abscess in the bed of the gallbladder | 2 (0,1%)                   |
| Suppuration of the epigastric puncture           | 12 (0,6%)                  |
| Total complications                              | 22 (1,1%)8                 |

Laparoscopic interventions made it possible to reduce the length of stay of patients in the hospital (by 22.2 - 37.5%), reduce the percentage of unnecessary laparotomies (by 15-25%), shorten the time of diagnosis, eliminate the occurrence of postoperative hernias, reduce the appointment of narcotic analgesics, transfer patients to enteral nutrition at an earlier date.

**Conclusion:** Within the framework of general scientific and technological progress, the constant introduction of new technologies makes them an integral part of a particular field of medicine. The current trend in the development of surgery is to reduce the invasiveness and traumaticity of surgical interventions. Due to the development of surgical technologies, one of the main tasks in surgery is to minimize surgical trauma and the subsequent reduction in the number of postoperative complications and mortality, as well as the duration of inpatient treatment of patients while maintaining the quality of surgical care. Today, urgent surgery can no longer be imagined without laparoscopy. The decision is more difficult in patients with unresectable disease who may be poor operative risk. The results show that surgical and endoscopic approaches have both advantages and disadvantages. We did not address quality of life issues directly and did not attempt to assess the relative cost-effectiveness of the procedures under study. These items, as well as the level of specialistic expertise available, will influence the therapeutic decision, that should be taken by the attending physician in strict cooperation with the fully informed patient.

#### List of literature

1. Abdullaeva, D. T., & Kh, A. G. (2023). Questionnaire for primary care physicians in Tashkent on connective tissue dysplasia.
2. Abdullaeva, D. T., Agzamkhodzhaeva, N. S., & Satibaldieva, N. R. (2023). COMBINATION OF IRRITABLE BOWEL SYNDROME AND BRONCHIAL ASTHMA ON THE BACKGROUND OF CONNECTIVE TISSUE DYSPLASIA IN CHILDREN. *Open Access Repository*, 9(1), 66-69.
3. Абдуллаева, Д. Т., Курбанова, Д. Р., & Гиёсова, У. Т. (2017). Нарушение психоэмоционального статуса у детей с бронхиальной астмой. *Научная дискуссия: вопросы медицины*, (1), 6-12.
4. Абдуллаева, Д. Т. (2011). Значение дисплазии соединительной ткани в развитии и прогнозе бронхиальной астмы у детей.
5. Халматова, Б. Т., Сотиболдиева, Н. Р., & Абдуллаева, Д. Т. (2010). Отдаленные последствия бронхообструктивного синдрома у детей. *Врач-аспирант*, 43(6), 57-61.



6. Khodadadi, H., Taghizadeh, M., Shabankareh, F., & Pakshir, K. (2021). Climate changes and emerging fungal infections. *Tehran University Medical Journal TUMS Publications*, 78(10), 684-693.
7. Xabibullayevna, M. M., TELMANOVNA, A. D., GAYRATOVNA, A. D., ABDUKARIMOVNA, S. M., & MASRUROVNA, A. M. (2021). Antileukotriene Drugs in The Treatment of Atopic Dermatitis in Children. *International Journal of Pharmaceutical Research*, 13(1), 2117-2121.
8. Mamatmusaeva, F., Tuychiev, L., Nuruzova, Z., Yodgorova, N., & Orinbaeva, Z. (2020). Optimizing the treatment of biliary disease in children with viral hepatitis. *International Journal of Pharmaceutical Research*, 12(4), 536-541.
9. Даминов, Т. О., Туйчиев, Л. Н., Худайкулова, Г. К., Маматмусаева, Ф. Ш., Аладова, Л. Ю., & Собирова, Г. Н. (2014). Нарушение моторной функции билиарной системы у реконвалесцентов вирусных гепатитов А и В и методы ее коррекции. *Детские инфекции*, 13(2), 16-19.
10. Юсупалиева, Г. А., & Иноятова, Ф. И. (2017). Возможности комплексных эхографических исследований в диагностике хронических вирусных гепатитов у детей. *Журнал теоретической и клинической медицины*, (1), 107-110.
11. Маматмусаева, Ф. Ш., Джураева, З. Б., Оринбаева, З. Н., & Юлдошева, Н. Г. (2022). ВИРУСЛИ ГЕПАТИТ «С» БИЛАН КАСАЛЛАНГАН БОЛАЛАРДА БИЛИАР ТИЗИМ ЎЗГАРИШЛАРИНИНГ БИОКИМЁВИЙ ХУСУСИЯТЛАРИ.
12. Муминова, М. Т., & Маматмусаева, Ф. Ш. (2022). Ўткир диареяли ОИВ зарарланган болаларда ичакнинг факультатив микрофлорасига *Sacharomyces bouiladining* таъсири.
13. Туйчиев, Л. Н. (2019). Вирусли гепатит С билан касалланган болаларда билиар тизим ўзгаришларининг клиник биокимёвий-усуллари.
14. Даминов, Т. О., Туйчиев, Л. Н., Худайкулова, Г. К., Собирова, Г. Н., Муминова, М. Т., & Маматмусаева, Ф. Ш. (2012). Биохимический состав желчи у реконвалесцентов гепатита А. *Детские инфекции*, 11(4), 57-60.
15. Ermatova, G. A., Teshaboev, U., & Mamatmusayeva, G. T. F. S. (2022). *MOTORIC FUNCTION OF THE BILIARY SYSTEM AT CHILDREN CONVALESCENTIA VIRAL HEPATITIS" A" AND" B" AND THEIR CORRECTION* (Doctoral dissertation).
16. Исмоилова, М. И. (2022). РОЛЬ КАЛЬПРОТЕКТИНА В ДИАГНОСТИКЕ ВОСПАЛИТЕЛЬНЫХ ПРОЦЕССОВ В КИШЕЧНИКЕ У БОЛЬНЫХ COVID-19. *Университетский терапевтический вестник*, 4, 62-62.
17. Ismailova, M. I., & Gadaev, A. G. (2022). Studying and Improving Treatment of Calprotectin, Helicobacter Pylori and Interleukin-6 in Blood of Patients with Covid-19. *Journal of Pharmaceutical Negative Results*, 2387-2393.
18. Gadaev, A., Ismoilova, M., & Turakulov, R. (2022). Comparative analysis of calprotectin and helicobacter pylori in the faces and interleukin-6 in the blood of patients with and without COVID-19 before and after the treatment. *Scientific Collection «InterConf+»*, (26 (129)), 236-242.
19. Ismoilova, M. I. (2022). Comparative Analysis of Calprotectin and Helicobacter Pylori in the Faces and Interleukin-6 in the Blood of Patients with and without Covid-19 Before and After the Treatment. *Central Asian Journal of Medical and Natural Science*, 3(5), 218-222.
20. Исмоилова, М. И., & Гадаев, А. Г. (2022). COVID-19 ўтказган ва ўтказмаган ошқозон-ичак тизимида патологик ўзгаришлар аниқланган беморларда ичаклар дисбактериозини аниқлаш.
21. Ismoilova, M. I., & Gadayev, A. G. (2022). *Comparative Study of Calprotectin in Feces and Interleukin-6 in the Blood of Patients with Covid-19* (Doctoral dissertation, America).

22. Гадаев, А. Г., & Исмоилова, М. И. (2022). COVID-19 ўтказган беморлар нажасида кальпротектин, helicobacter pylori ва конида интерлейкин-6 кўрсаткичларининг солиштирма тахлили.
23. Mahammadovna, S. I. (2023). Features of Cluster Design in Modern Paradigms of Education. *Telematique*, 22(01), 348-355.
24. Сирожиддинова, И. (2022). Методика смешанной отборки при комплексном проектировании профессиональной подготовки будущих инженеров. *Общество и инновации*, 3(7/S), 87-92.
25. Rakhmonberdiyevna, T. S. (2022). CREATIVITY AS A PEDAGOGICAL PROBLEM. *Conferencea*, 138-141.
26. Mahammadovna, S. I. (2022). IMPROVING THE PROFESSIONAL TRAINING OF FUTURE ENGINEERS BASED ON THE CLUSTER APPROACH. *Spectrum Journal of Innovation, Reforms and Development*, 3, 45-47.
27. Sirojiddinova, I. M. (2015). ENGINEERING STUDENTS HAVE SUCCEEDED IN CREATING A TECHNOLOGY CLUSTER. *Pedagogy & Psychology. Theory and practice*, 22.
28. Sirojiddinova, I. (2023). TECHNOLOGICAL CHARACTER OF THE EDUCATIONAL PROCESS WHEN DESIGNING PEDAGOGICAL OBJECTS. *Solution of social problems in management and economy*, 2(2), 130-132.
29. Sirojiddinova, I. M. (2015). ENGINEERING STUDENTS HAVE SUCCEEDED IN CREATING A TECHNOLOGY CLUSTER. *Pedagogy & Psychology. Theory and practice*, 22.
30. Mahammadovna, S. I. (2022, October). DEVELOPMENT OF A METHODOLOGICAL SYSTEM OF TRAINING BASED ON THE CLUSTER APPROACH. In *Archive of Conferences* (pp. 30-33).
31. Сирожиддинова, И. (2022). Методика смешанной отборки при комплексном проектировании профессиональной подготовки будущих инженеров. *Общество и инновации*, 3(7/S), 87-92.
32. Mahammadovna, S. I. (2021). Needs and factors for developing professional and creative abilities of students of higher educational institutions. *Annals of the Romanian Society for Cell Biology*, 25(6), 2200-2209.
33. Медведев, С. Е., Волкова, В. Н., Волков, Н. Н., & Чухловина, М. Л. (2018). Особенности церебральной патологии у беременных: результаты ретроспективного одномоментного исследования. *Фарматека*, (6), 359.
34. Карлов, В. А., Куликов, Ю. А., Ильина, Н. Л., & Грабовская, Н. В. (1997). Дисциркуляторная энцефалопатия у больных артериальной гипертензией. *Журнал неврологии и психиатрии им. СС Корсакова*, 97(5), 15-17.
35. Мамурова, М. М., Джурабекова, А. Т., & Игамова, С. С. (2021). Оценка когнитивных вызванных потенциалов головного мозга (p-300) у лиц молодого возраста с артериальной гипотензией. *журнал неврологии и нейрохирургических исследований*, 2(1).
36. Salimovna, S. D., Suratovna, I. S., Tahirovna, D. A., Vladimirovna, S. K., Fedorovna, V. N., & Hamzayevna, M. M. (2019). Endothelial dysfunction factor as an indicator of clinical and neurological disorders in pregnant women with preeclampsia. *Достижения науки и образования*, (11 (52)), 55-60.
37. Takhirovna, D. A., Otabekovich, S. A., Axmatjonovich, G. A., & Mirxamzaevna, M. M. (2021). The Nature Of Cognitive Impairment In Patients With Astheno-Neurotic Syndrome. *nveo-natural volatiles & essential oils journal/ nveo*, 5942-5948.
38. Ergasheva, M., & Vakhobova, N. (2019). New gender-influenced stroke study: Cognitive manifestations in acute ischemic stroke in Uzbekistan. *Journal of the Neurological Sciences*, 405, 115.

39. Ergasheva, M., Vakhobova, N., & Rakhimbaeva, G. (2019). Gender, aging and background diseases influence on the new neuronosological structure of acute ischemic stroke in Uzbekistan. *Journal of the Neurological Sciences*, 405, 115.
40. Tolibova, N., & Vakhobova, N. (2017). Gender differences in stroke subtypes, severity, risk factors, and outcomes among elderly patients with acute ischemic stroke in Uzbekistan. *Journal of the Neurological Sciences*, 381, 377.
41. Makhmudovich, A. M., Sattarovna, R. G., Maksudovna, V. N., & Azamatovich, J. S. (2021). Hyperhomocysteinemia And Pathogenetic Mechanisms Of Ischemic Stroke. *The American Journal of Medical Sciences and Pharmaceutical Research*, 3(02), 66-76.
42. Tolibova, N., & Vakhobova, N. (2017). Stroke incidence and association with risk factors in women in Uzbekistan. *Journal of the Neurological Sciences*, 381, 377.
43. Асадуллаев, М. М., Саидвалиев, Ф. С., Шермухамедова, Ф. К., Ризвонов, Ж. К., & Вахабова, Н. М. (2012). Оценка мультимодального действия цитофлавина при остром мозговом инсульте, развившемся на фоне метаболического синдрома. *Журнал неврологии и психиатрии им. СС Корсакова*, 112(10), 24-27.
44. Makhmudova, U., Rakhimbaeva, G., & Vakhobova, N. (2019). New approach of risk factors and background diseases role in acute ischemic stroke in elderly and senile-aged patients in Uzbekistan. *Journal of the Neurological Sciences*, 405, 118.
45. Allaberganov Azizbek Saparbeyevich. (2023). SHIFOKOR – TIBBIYOT XODIMLARI – BEMOR MUNOSABATLARINING XUSUSIYATLARI. AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI, 2(2), 208–211. Retrieved from <https://sciencebox.uz/index.php/amaltibbiyot/article/view/5963>
46. Allaberganov Azizbek Saparbeyevich, Safarniyozova Dinora, Alimova Komila, & Rasulov Ma'murjon. (2023). FUQAROLARNING TIBBIY IJTIMOIIY YORDAM Olish HUQUQI. AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI, 2(2), 223–227. Retrieved from <https://sciencebox.uz/index.php/amaltibbiyot/article/view/5966>
47. Эркинов Жахонгир Хайрулла Угли, & Аллаберганов Азизбек Сапарбоевич (2023). МЕСТО СОЦИОЛОГИИ МЕДИЦИНЫ В СИСТЕМЕ СОЦИАЛЬНОГО ЗНАНИЯ. Ta'lim fidoyilari, 14-02 , 16-18.
48. Toxirova Nigina Toxirovna, & Аллаберганов Азизбек Сапарбоевич (2023). DEVIANT XULQ-ATVOR SOTSIOLOGIYASI. Ta'lim fidoyilari, 14-02 , 12-15.
49. Ш. С. Шаниязов, & А. С. Аллаберганов (2022). ЎЗБЕКИСТОНДА БИОЭТИКАНИНГ ВУЖУДГА КЕЛИШИ ВА РИВОЖЛАНИШИ ТАРИХИДАН. Academic research in educational sciences, TSDI and TMA Conference (2), 95-97.
50. Исмоилова, М. И. (2022). РОЛЬ КАЛЬПРОТЕКТИНА В ДИАГНОСТИКЕ ВОСПАЛИТЕЛЬНЫХ ПРОЦЕССОВ В КИШЕЧНИКЕ У БОЛЬНЫХ COVID-19. *Университетский терапевтический вестник*, 4, 62-62.
51. Ismailova, M. I., & Gadaev, A. G. (2022). Studying and Improving Treatment of Calprotectin, Helicobacter Pylori and Interleukin-6 in Blood of Patients with Covid-19. *Journal of Pharmaceutical Negative Results*, 2387-2393.
52. Gadaev, A., Ismoilova, M., & Turakulov, R. (2022). Comparative analysis of calprotectin and helicobacter pylori in the faces and interleukin-6 in the blood of patients with and without COVID-19 before and after the treatment. *Scientific Collection «InterConf+»*, (26 (129)), 236-242.
53. Gadaev, A., Ismoilova, M., & Turakulov, R. (2022). Comparative analysis of calprotectin and helicobacter pylori in the faces and interleukin-6 in the blood of patients with and without COVID-19 before and after the treatment. *Scientific Collection «InterConf+»*, (26 (129)), 236-242.

54. Исмоилова, М. И., & Гадаев, А. Г. (2022). COVID-19 ўтказган ва ўтказмаган ошқозон-ичак тизимида патологик ўзгаришлар аниқланган беморларда ичаклар дисбактериозини аниқлаш.
55. Ismoilova, M. I., & Gadayev, A. G. (2022). *Comparative Study of Calprotectin in Feces and Interleukin-6 in the Blood of Patients with Covid-19* (Doctoral dissertation, America).
56. Гадаев, А. Г., & Исмоилова, М. И. (2022). COVID-19 ўтказган беморлар нажасида кальпротектин, helicobacter pylori ва конида интерлейкин-6 кўрсаткичларининг солиштирма таҳлили.
57. Isroilovna, I. M., & O'G'Li, I. O. B. (2021). KATTALARDAGI ONKOLOGIK KASALLIKLARNI PSIXOLOGIK TAHLILI. *Oriental renaissance: Innovative, educational, natural and social sciences*, 1(4), 516-522.
58. Ismoilova, M. I., & Gadayev, A. G. (2022). *Comparative Study of Calprotectin in Feces and Interleukin-6 in the Blood of Patients with Covid-19* (Doctoral dissertation, America).