

### Application of Transpedicular Fixers in Surgery Treatment of Spondylolisthesis

**B. M. Isakov**

Andijan State Medical Institute

#### Article Information

**Received:** June 21, 2023

**Accepted:** July 22, 2023

**Published:** Aug 23, 2023

**Keywords:** *surgical treatment, extensor muscles, spinal fusion, lumbosacral region.*

#### ABSTRACT

The aim of this work is to analyze our own results of surgical treatment of 30 patients with isthmic spondylolisthesis of the L5 vertebra, who used the method of transpedicular fixation (TPF) using various types of transpedicular constructs and methods of their installation in combination with various methods of anterior fusion.

The results of surgical treatment of 30 patients with isthmic spondylolisthesis of the L5 vertebra were analyzed. Displacements of I–II degree were present in 10 patients, III–IV degree – in 17 patients, and spondyloptosis – in 3 patients. Depending on the level of fixation, 4 groups were distinguished: patients of group I received fixation of two segments, screws were inserted into L4, L5, and S1 vertebrae; in the remaining groups, three segments were fixed, while in group II, screws were inserted into L3, L4, L5, and S1 vertebrae, in group III, into L3, L5, and S1 vertebrae, and in group IV, into L3, L4, and S1 vertebrae. Posterior fixation was combined with L5–S1 interbody fusion performed using various techniques.

#### Clinical results of treatment

were regarded in 93.3% of cases as good. Satisfactory results in 6.7% of cases occurred due to the development of mild paresis of the extensor muscles of the foot and fingers in two patients after surgery. Long-term results were followed up in 16 patients, the average follow-up period was 29 months. Radiologically, with the exception of patients with structural fractures, fixation of the lumbosacral region was stable. Fracture of metal structures occurred in 7 patients (23.3%), 6 of whom did not undergo anterior spinal fusion. In this case, there was a partial loss of reduction. Clinically, metal structure fractures did not lead to deterioration. After L5–S1 interbody fusion with cortical grafts, stabilization of the lumbosacral region also occurred. The calculation of radiological parameters characterizing the change in the anatomical and biomechanical relationships of the lumbosacral region before and after surgical treatment was carried out. Analysis of the results of treatment indicates the effectiveness of TPF in combination with interbody fusion in the treatment of spondylolisthesis. The use of L5–S1 cortical graft in severe degrees of spondylolisthesis has been substantiated. The goal of surgical treatment for

spondylolisthesis is to eliminate pain, restore anatomical relationships and stabilize the lumbosacral spine [16]. The technique of transpedicular fixation, which is widely used in the surgical treatment of spinal pathology, including spondylolisthesis, has shown its high efficiency [6]. The importance of L5–S1 interbody fusion is emphasized by most authors, and a variety of methods and techniques for its implementation are also reported [2–4, 11, 12, 16]. There are no data in the literature on a comparative analysis of the results for various methods of installing pedicle fixators.

### **Material and methods**

57 patients with spondylolisthesis underwent surgical treatment using transpedicular fixation in the Department of Neurosurgery of the AF RRCEMMP. In most cases (30 patients) there was isthmic spondylolisthesis of the L5 vertebra. We analyzed the treatment of this group of patients. The age distribution was as follows: from 18 to 40 years - 17 and from 41 to 55 years - 13. Male - 18 patients, female - 12. In the complex of preoperative instrumental examination, patients underwent standard and functional radiography of the lumbosacral spine and myelography, CT combined with myelography and MRI. In 22 patients, a spondylolysis defect of the L5 vertebral arch was detected. In 8 cases, such dysplastic changes were observed as a flattened elongated arch of the L5 vertebra, a rounded deformed upper endplate of the first sacral vertebra, and in 6 cases, non-fusion of the posterior elements of the lumbar and sacral vertebrae. Depending on the degree of displacement according to Meyerding, the distribution of patients was as follows: spondylolisthesis of I degree - in two, II degree - in eight, III degree - in twelve, IV degree - in five; three patients had spondyloptosis. Thus, displacements of I-II degree in 33.3% of patients, severe and extremely severe - in 66.7%. Patients in all cases complained of pain, leading to limitation of physical activity. Complaints of pain only in the lumbosacral region were presented by 5 patients, pain both in the lumbosacral region and in the lower extremities was disturbed by 25 patients, and in 6 of them the irradiation of pain was bilateral. In one 34-year-old patient, lumbodynia was combined with the syndrome of intermittent myelogenous claudication. In one case, when visiting a 39-year-old patient with spondylitis) interlink angle S1 - between the lines connecting the centers of the body L5–S1 and S1–S2.

### **Results**

Clinically, in the postoperative period and during further follow-up, all patients showed complete regression of pain symptoms. All patients returned to full-fledged physical activity within a year after the operation, continued their educational and labor activities. All patients, with the exception of two who developed neurological complications after surgery, were completely satisfied with the treatment.

### **Discussion**

The issues of surgical treatment of spondylolisthesis are an actual problem of vertebrology and are widely discussed in modern literature. As noted by almost all authors, there is currently no unity in the choice of tactics and methods of surgical treatment of spondylolisthesis. The method of transpedicular fixation, which is widely used in modern spinal surgery, including the treatment of spondylolisthesis, has a number of advantages over others. This is the possibility of segmental fixation, rigidity and primary stability of fixation, which can significantly reduce the time of functional recovery after surgery. The improvement of transpedicular fixators and the use of multiaxial screws facilitates the installation of the structure in case of deformities and makes it possible to reduce the displaced vertebra from the posterior approach [17]. Particular importance is attached to the issue of reduction of the displaced vertebra and elimination of anatomical and biomechanical disorders occurring in spondylolisthesis. To assess these changes and their correction after surgery, both our scientists and foreign scientists proposed a number of radiometric characteristics [1, 9, 18]. If, with small degrees of displacement, a pronounced

reduction is possible, then in cases of large displacements, the reduction is much more complicated and to a certain extent limited. When using reduction and fixation with transpedicular constructs for severe displacements, in general, our results in a number of parameters are comparable with the literature data [11, 16], except for the cases when the reduction was performed by anterior resection of the displaced vertebra and fixation of L4 to S1. The reduction rates in such cases are higher, but it is necessary to note the great technical difficulties and risks associated with such operations [3, 16].

### **References:**

1. Antonov I.P. Current state and prospects for studying diseases of the peripheral nervous system. // VIII All-Union Congress of neuropathologists, psychiatrists and narcologists. - M.: 1988. - v. 3. -
2. Antonov I.D., Nezdved G.K., Ivashina E.N. and others. To the pathogenesis of neurological manifestations of lumbar osteochondrosis / Clinic, treatment and prevention of diseases of the nervous system: Materials of the republican scientific-practical conference. - Kazan: 1988. - S. 65-70.
3. Arutyunov A.I., Brotman M.K. Contrast research methods in the clinical diagnosis of degenerative processes in intervertebral discs // Questions of Surgery, 1964. - No. 5. - S. 44-49.
4. Blagodatsky M.D., Meyerovich S.I. Diagnosis and treatment of discogenic lumbosacral sciatica. - Irkutsk: Publishing House of Irkutsk University, 1987. - 272 p.
5. Brotman M.K. Neurological manifestations of lumbar osteochondrosis. - Kyiv: Health, 1975. - 168 p.
6. Bugakchiyskaya N.M. Manual therapy in the complex treatment of neurological manifestations of lumbar osteochondrosis: Ph.D. diss. ... cand. med.n. - Kyiv: 1987. - 21 p.
7. Veselovsky V.P. Practical vertebroneurology and manual therapy. - Riga, 1991. - 341 p.
8. Gekht B.M., Kasatkina L.F., Samoilov M.I., Sanadze A.G. Electromyography in the diagnosis of neuromuscular diseases. - Taganrog: TRTU, 1997. - 370 p.
9. Golubkov O.I. Outcome and recovery of working capacity of patients after conservative and surgical methods of treatment of lumbar osteochondrosis: author. diss. ... cand. med.n. - M.: 1972. - 17 p.
10. Gusev E.I., Konovalov A.N., Belyakov V.V. etc. Research methods in neurology and neurosurgery. - M.: Knowledge, 2000. - 336 p.
11. Zharkov P.L. Osteochondrosis and other changes in the spine in adults and children. - M.: Medicine, 1994. - S. 23-35.
12. Zhulev N.M., Badzgaradze Yu.D., Zhulev S.N. Osteocondritis of the spine. - St. Petersburg: 2001. - 588 p.
13. Ivanichev G.A. Clinical and electromyographic assessment of the coordination relations of the leg muscles in lumbar osteochondrosis. / Modern methods of research in neurology and psychiatry. - Kursk: 1977. - v. 2. - S. 109-111.
14. Kazmin A.I., Kaptelin A.F. Indications for conservative orthopedic treatment in discogenic pain syndrome. - M.: 1974. -
15. Kazmin A.I. Operative treatment of lumbar osteochondrosis. // Orthopedic traumatology, 1986. - No. 10. - S. 17-20.

16. Kinzersky A.Yu. Dopplerography of the epidural venous plexuses in differentiating the causes of compression of the spinal cord roots in intervertebral osteochondrosis. // Visualization in the clinic, 1999. - No. 14-15. - S. 44-46.
17. Kipervas I.P. Peripheral neurovascular syndromes. - M.: Medicine, 1985. - 175 p.
18. Musalatov Kh.A., Aganesov A.G. Surgical rehabilitation of radicular syndrome in osteochondrosis of the lumbar spine (microsurgical and puncture discectomy). - M.: Medicine, 1998. - 88 p.