SCHOLASTIC:

Journal of Natural and Medical Education

Volume 2 Issue 2, Year 2023 ISSN: 2835-303X https://univerpubl.com/index.php/scholastic

Taxonomic Indicators of Congenital Heart Defects and their Prevalence in Hereditary Diseases

Khudoykulova Nasiba Ismailovna

Bukhara State Medical Institute

Article Information

Received: December 09, 2022 Accepted: January 10, 2023 Published: February 11, 2023

Keywords: Down syndrome, anomaly, congenital anomales, polydactyly, syndactyly.

ABSTRACT

Carried out an epidemiological study on the prevalence of congenital anomalies of malformations, (CAM) in Bukhararegion for the period from 2019 to 2021. It was shown that these CAMs as congenital heart defect and Down's syndrome were the most common pathology among other CAMs.

It is well known that. The Bukhara region is an important economically developed industrial region located in the arid zone of Uzbekistan, which is characterized by the presence of a sharply continental climate, a high level of insolation, a high degree of technogenic impact caused by the operation of industrial enterprises. The complex of technogenic factors affecting the health of the local population can be attributed to the so-called "controlled risk factors", on the basis of which it seems promising to develop and implement a target program for the prevention and treatment of congenital malformations congenital anomalesin the region [3,6]. In recent years, a continuous growth of VAR has been detected in the world. An increase in the relative indicators of hereditary and congenital pathology in the structure of morbidity and mortality of newborns and young children was noted [1,7]. It is shown that more than 10% of all conceptions in the human population are accompanied by congenital anomales of which 0.5% are chromosomal diseases, 0.7% are molecular, 1.8% are polygenic pathologies, the remaining 7% are hereditary predispositions [I, 7]. 4-6% of children with congenital anomalesare born annually in the world, and in many countries, fetal and newborn congenital anomales are among the first in the structure of mortality [1,4, 7]. In connection with the above, the study of the frequency of spread, early diagnosis, prevention and treatment of congenital anomalesis one of the urgent tasks of modern medicine.

The purpose of the work is to analyze the frequency of distribution of a number of VAR in the Bukhara region.

Material and methods of research. We conducted prospective epidemiological studies that included detailed registration of congenital anomales in the Bukhara region. It should be noted that only based on accurate data on the frequency of occurrence of congenital anomales in the

regions, and then in Uzbekistan, it is possible in the future to judge the epidemiological picture and the frequency of occurrence of congenital anomales, to conduct a medical examination of children with congenital anomalesand determine the required amount of pediatric and other specialized types of treatment [4, 5].

Individual examination cards were developed for each child with congenital anomales with the introduction of objective and reliable information there, starting from the beginning of the mother's pregnancy, the development of the child - to the profiling clinics, taking into account the diagnoses.

In this study, archival materials of maternity hospitals in the city of Bukhara were used .- The Bukhara region, as well as the data of the Bukhara Regional Screening Center. In maternity hospitals, all the birth histories and the history of the development of newborns for the period from 2019 to 2021 were studied and analyzed.

Results and discussion. The results of the research showed that 84,159 live children were registered in the settlements/ Bukhara region for 4 years.

The prevalence rates of 6 types of congenital anomaleswere analyzed (Table). Of these, the following types of congenital anomales were most common: congenital heart defects, Down syndrome, spinal hernia and polydactyly.

For example, the frequency of congenital heart defects was 0.48 or

1:2053, with Down syndrome - 0.40 or 1 case per 2475 newborns, with spinal hernia - 0.34 or 1 : 2902, with polydactyly - 0.32 or 1 case per 3117.

Tablel. Quantitative characteristics of congenital anomales in the Bukhara region for the period from 2019 to 2021

.Vo	Name of BA	Theamount of INJECTED				
		2019	2020	2021	Total	X
I.	Congenitaltorticollis	10	6	7	23	0,27 И1 : 3659
2.	DownSyndrome	14	12	8	34	0,40 // 1 : 2475
3.	Congenitalheartdefects	14	12	14	40	0,48 <i>И</i> 1 : 2053
4.	Polydactyly	14	7	6	27	0,32//1:3117
5.	Syndactyly	3	4	5	12	0,14//1:7013
6.	Spinalcоггрыжа	10	9	10	29	0,34 <i>U</i> 1 : 2902

I Note: X is the frequency per 1000 live births over 4 years.

The second group of VPD, namely congenital torticollis and syndactyly, occurred with a lower frequency compared to the above-mentioned congenital anomales and according to the frequency of epidemiological prevalence in the Bukhara region, they can be conditionally attributed to the group of pathologies with a moderate nature of the spread of congenital anomales ..

Congenital torticollis was registered with a frequency of 0.27 or 1 case per 3659 newborns. Syndactyly in terms of the frequency of occurrence of congenital anomales was in last place: 0.14 or 1: 7013.

The results obtained indicate that the epidemiological picture of the situation in the Bukhara region is unfavorable. We obtained different frequencies of congenital anomales propagation.

Higher frequencies of the epidemiological prevalence of congenital anomales were recorded here, in contrast to those in the Navoi region [2].

It should be emphasized that in the Bukhara region, unfavorable environmental impacts in the complex negatively affect the health of the population living here [3, 6] and make a certain

^{© 2023} by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/).

"contribution" to the frequency of birth of children with congenital anomales.

We believe that the high frequency of congenital anomalesoccurrence in this region is due to environmental, hereditary and social factors.

CONCLUSIONS:

- 1. We have obtained epidemiological data on the general picture of the frequency of congenital anomalesspread in the Bukhara region.
- 2. In the Bukhara region, a steady dynamic growth of congenital anomaleswas noted.
- 3. In this region, a high incidence of congenital heart defects, Down syndrome, spinal hernia and polydactyly was recorded.
- 4. Apparently, a complex of unfavorable environmental factors in the Bukhara region

The region affects the epidemiological situation of the frequency of the spread of VAR.

REFERENCES

- 1. Худойкулова, Н.И. (2022). АНАЛИЗ И РАСПРОСТРАНЕНИЕ ВРОЖДЕННЫХ АНОМАЛИЙ В БУХАРСКОЙ ОБЛАСТИ. Scientific progress, 3(1), 954-957.
- 2. Худойкулова Н.И. (2018). Пути воспитания толерантности у молодежи. *Наука, техника и образование*, (11 (52)), 98-100.
- 3. Худайкулова, Н.И. (2022). ХИМИЧЕСКИЕ И ФИЗИЧЕСКИЕ ФАКТОРЫ И ИХ ВЛИЯНИЕ НА ИММУННУЮ СИСТЕМУ ОРГАНИЗМА. *Scientificprogress*, *3*(1), 891-895.
- 4. KHUDOYKULOVA N.I. (2022, February). CONGENITAL ANOMALIES, ANALYSIS AND DEVELOPMENT IN THE NAVOI REGION. *INTERNATIONAL JOURNAL OF PHILOSOPHICAL STUDIES AND SOCIAL SCIENCES* 2(2) (pp.46-49).
- 5. Худойкулова, Н. И. (2022). ПУТИВ НЕДРЕНИЯ ОХОТНИЧЬЕГО ТУРИЗМА В ОХОТНИЧЬИХ ХОЗЯЙСТВАХ . International Journal of Philosophical Studies and Social Sciences, 2(2), 213-219.
- 6. Khudaikulova N. I. (2022). EXTERNAL FACTORS AND THEIR INFLUENCE ON THE BODY'S IMMUNE SYSTEM .(pp.343-348)
- 7. Н.И. Худойкулова, М.А.Усманова (март, 2022).ЭКОЛОГИЧЕСКИЕ ОСОБЕННОСТИ ЖИВОТНЫХ, ОБИТАЕМЫХ В ПУСТЫННОЙ ЗОНЕ БУХАРСКОЙ ОБЛАСТИ Н.И. Международный научно-образовательный электронный журнал «ОБРАЗОВАНИЕ И НАУКА В XXI ВЕКЕ». Выпуск №24 (том 6) стр.1359-1362
- 8. Khudaikulova N.I. (2022) FACTORS AFFECTING THE HUMAN IMMUNE SYSTEM Международный научно-образовательный электронный журнал «ОБРАЗОВАНИЕ И НАУКА В XXI ВЕКЕ». Выпуск №24 (том 6) (март, 2022). Стр.1363-1367.
- 9. Худойкулова Н.И.(2021)Ўзбекистоннинг чўл зонасида овланадиган ҳайвон турлари ҳақида янги маълумотлар Хоразм Маъмун академияси ахборотномаси: илмий журнал.-№11 (82.) 75-78 б.

- 10. *Худайкулова Н.И., Назарова Ф.И.*(2021) Конкурентоспособность и качество продукции в сфере текстильной промышленности Хоразм Маъмун академияси ахборотномаси: илмий журнал.-№12 (83), Хоразм Маъмун академияси, 158 -161 б.
- 11. *Худойкулова Н.И.* (2020) Тўдакўл сув омбори зоопланктонлари ҳақида маълумот Хоразм Маъмун академияси ахборотномаси: илмий журнал.-№9(66), Хоразм Маъмун академияси, 68-70 б.
- 12. Ilhomovna, F. N. (2022). Let" Wolf Mouths" and" Hares" Not Be Born.... *Research Journal of Trauma and Disability Studies*, *1*(12), 38-44.
- 13. Ilkhomovna, N. F. (2022). Negative Impact of Seizures on Quality of Life. *Miasto Przyszłości*, 24, 120-122.
- 14. Nazarova, F. I. (2022). SOG'LOM FARZAND OILA QUVONCHI. *Scientific progress*, *3*(2), 1010-1015.
- 15. Ilhomovna, F. N. (2022). LATE SEIZURES AND CONSEQUENCES OF EPILEPSY IN YOUNG CHILDREN. *Conferencea*, 219-223.
- 16. Nazarova, F. (2022). Qaridoshlar orasidagi ofat. Scientific progress, 3(1), 663-669.
- 17. Nazarova, F. I. (2022). ABU ALI IBN SINONING SOGʻLOM TURMUSH TARZINI SHAKILANIRISHI HAQIDA. *Scientific progress*, *3*(1), 1137-1142.