

### Verification, Typing and Study of Sensitivity of N.Gonorrhoeae to Antimicrobial Drugs is Carried out Using a Complex of Microbiological and Molecular Biological Methods

**Munisa Khamrakulovna Ablakulova**

General Practice of Family Medicine, Samarkand State Medical University

**Khusinova Shoira Akbarovna**

Candidate of Medical Sciences, Associate Professor, Samarkand State Medical University,  
Samarkand, Uzbekistan

#### Article Information

**Received:** November 29, 2022

**Accepted:** December 30, 2022

**Published:** January 31, 2023

**Keywords:** *n. gonorrhoeae*, sensitive, typing and research.

#### ABSTRACT

*this article attempts to reveal the main reasons for the use of antibiotic-resistant strains of the causative agent of gonococcal infection as one of the directions for the formation of a national collection of pathogenic microorganisms. To carry out scientific work, the author conducted an experiment on where viable strains of N.gonorrhoeae obtained as part of the program for monitoring the antibiotic resistance of STI pathogens in the territory. Verification, typing and sensitivity testing of N.gonorrhoeae to antimicrobial agents is carried out using a complex of microbiological and molecular biological methods. The problem in question is still little studied, therefore, requires more thorough research.*

**Introduction:** Gonococcus (lat. Neisseria gonorrhoeae) is a species of Gram-negative diplococci of the genus Neisseria. Cause gonorrhea - anthroponotic venereal infection, characterized by purulent inflammation of the mucous membranes, often the genitourinary system. Transmission of pathogens occurs, as a rule, sexually or through personal belongings. In 10% of infected men and 80% of infected women, the disease is asymptomatic.

Due to the common transmission routes of the pathogen, a mixed infection is possible, in which, in addition to gonococci, there are chlamydia, ureaplasma, Trichomonas, Candida fungi, herpes viruses, etc. It is known that gonorrhea with concomitant trichomoniasis is treated with great difficulty, and this is due to the fact that trichomonads - the causative agents of trichomoniasis - "absorb" gonococci and make them inaccessible to antibiotics. In addition, gonococci are capable of endocytobiosis - that is, to live and multiply inside other cells, in particular, inside blood cells - leukocytes. In such cases, after the death of gonococci, the inflammatory process can be supported by these microorganisms (post-gonorrheal diseases). Gonorrhea is often noted in patients with syphilis.

Gonococci parasitize mainly on mucous membranes lined with cylindrical or glandular epithelium (urethra, cervical canal, distal rectum, conjunctiva). Eye damage in adults occurs

due to the introduction of pathogens by hand from the genitourinary organs. It is also possible to infect girls at home in violation of hygienic rules ("potty" infection, sharing a bed with a sick gonorrhoea, using common personal hygiene items, such as sponges, etc.) Lesions of mucous membranes covered with stratified squamous epithelium, as well as skin lesions, rarely occur, only under special circumstances (trauma, hormonal changes in the body, immunodeficiency states). Reproducing on the surface of the epithelium, gonococci can cause its destruction and enter the superficial lymphatic and blood vessels. The time required for the penetration of gonococci into the subepithelial layer and the development of inflammation determines the duration of the incubation period: from 1-2 days to 1 month or more.

Diagnostic value has an increased content of leukocytes in the urine in the absence of any complaints. The diagnosis is considered proven only when pathogens are determined in smears or cultures

**Purpose:** deposition, storage and use of antibiotic-resistant strains of the causative agent of gonococcal infection as one of the directions for the formation of a national collection of pathogenic microorganisms

**Materials and methods:** viable strains of *N.gonorrhoeae* obtained as part of the program for monitoring the antibiotic resistance of STI pathogens in the territory. Verification, typing and sensitivity testing of *N.gonorrhoeae* to antimicrobial agents is carried out using a complex of microbiological and molecular biological methods. For the storage of biological samples, the cryopreservation method (-80 ° C) is used, the most valuable objects of storage are duplicated in a freeze-dried state. The electronic base of the collection is maintained using the specialized program "Diam: Bank of Microorganisms", adapted to the characteristics of the storage facilities.

**Results:** The main direction of their use is the analysis of long-term trends in changes in antibiotic resistance of gonococcal pathogens, allowing to update clinical guidelines for the management of patients with this STI. Characteristics of the collection stock in terms of sensitivity to antimicrobial drugs, as well as the presence of genetic determinants of antibiotic resistance, indicates a gradual restoration of the sensitivity of *N. gonorrhoeae* to penicillins, tetracyclines, and fluoroquinolones. A stable high level of sensitivity of *N. gonorrhoeae* to third-generation cephalosporins (primarily ceftriaxone) has been shown, which determines their importance as drugs of choice for the treatment of gonococcal infection.

An in-depth analysis of *N. gonorrhoeae* deposited in the collection using targeted and whole genome sequencing technologies allows us to assess the current stage of the molecular evolution of the causative agent of gonococcal infection, which consists in the preservation of a significant proportion of multi-resistant strains in the bacterial population, which are a potential resource for the emergence of epidemically dangerous genotypes. The characterized multidrug-resistant strains are a valuable element of the collection, potentially in demand when searching for reserve drugs for the treatment of gonococcal infection.

**Выводы:** *N. gonorrhoeae* can be considered as an element of the emerging national collection of pathogenic microorganisms, the use of which will allow assessing and preventing the risks of incurable forms of gonococcal infection.

#### **Literature:**

1. Хакимова Л. и др. Безопасное материнство и эффективный антенатальный уход //Общество и инновации. – 2021. – Т. 2. – №. 8/S. – С. 91-95.
2. Khakimova L. et al. Acne in Allergic Skin Diseases //Texas Journal of Medical Science. – 2022. – Т. 8. – С. 129-131.

3. Лапасов С. Х. и др. Диагностика, лечение и профилактика хронического гепатита с в условиях первичного звена медицинской помощи (обзор литературы) //Здобутки клінічної і експериментальної медицини. – 2017. – №. 2. – С. 13-21.
4. Akbarovna K. S. et al. ASSESSMENT OF THE PREVALENCE AND QUALITY OF CARE OF PATIENTS WITH HEART FAILURE IN PRIMARY CARE //Thematics Journal of Education. – 2022. – Т. 7. – №. 3.
5. Хакимова Л. Р. и др. Оценка эффективности обучения основам доказательной медицины в последипломном образовании врачей общей практики //Медицина и экология. – 2017. – №. 4 (85). – С. 130-133.
6. Akbarovna K. S. et al. REVIEW OF THE COURSE AND TREATMENT FEATURES OF COVID-19 PATIENTS WITH CONCOMITANT CARDIOVASCULAR DISEASE //Asian journal of pharmaceutical and biological research. – 2022. – Т. 11. – №. 2.
7. Аблакулова М., Абдухамидова Д. ОСОБЕННОСТИ СТАРЕЮЩЕГО ОРГАНИЗМА И ФАРМАКОТЕРАПИИ В ГЕРИАТРИИ //InterConf. – 2020.
8. Khakimova L. et al. acne in allergic skin diseases. texas journal of medical science, 8. – 1818.
9. Abdullaev X. D. et al. EVALUATION OF THE EFFECTIVENESS OF THE DRUG GEPON IN THE TREATMENT OF GENITAL CANDYLOMAS //Web of Scientist: International Scientific Research Journal. – 2021. – Т. 1. – №. 02. – С. 16-19.
10. Абдуллаев Х. Д. и др. Определение цитокинового статуса у больных с псориазом на фоне стандартного лечения //Общество и инновации. – 2020. – Т. 1. – №. 1/S. – С. 625-629.
11. Ахмедова М. М., Абдуллаев Х. Д., Камолова М. И. ОЦЕНКА ЭФФЕКТИВНОСТИ МЕТОДОВ ЛЕЧЕНИЯ ОНИХОМИКОЗОВ У ВЗРОСЛЫХ //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIIY JURNALI. – 2022. – Т. 2. – №. 3. – С. 186-190.
12. Тошев С. У., Аширов З. Ф., Абдуллаев Х. Д. ЛЕЧЕНИЕ ВИТИЛИГО 308-нм ЭКСИМЕРНЫМ ЛАЗЕРОМ //Актуальные аспекты медицинской деятельности. – 2021. – С. 240-243.
13. Davlatovich A. X., Xaydarjonovna X. S., Raximjon N. S. JYBYK BILAN BO'LGAN PROSTATITNI DAVOLASH USULINI TAKOMILLASHTIRISH //BARQARORLIK VA ETAKCHI TADQIQOTLAR ONLAYN ILMIIY JURNALI. – 2022. – С. 359-361.
14. Ахмедова М., Расулова Н., Абдуллаев Х. Изучение парциальных функций почек у детей раннего возраста с нефропатией обменного генеза //Журнал проблемы биологии и медицины. – 2016. – №. 2 (87). – С. 37-40.
15. Нарзикулов Р. и др. Принципы терапии у женщин больных гонореей ассоциированные иппп //Журнал вестник врача. – 2019. – Т. 1. – №. 1. – С. 99-102.
16. Нуруллаева А. А., Рахматова А. Х., Абдуллаев Х. Д. ЗНАЧЕНИЕ МИКРОБНОГО ОБСЕМЕНЕНИЯ КОЖИ ПРИ НЕКОТОРЫХ ЗУДЯЩИХ ДЕРМАТОЗАХ //Молодежь и медицинская наука в XXI веке. – 2019. – С. 125-125.
17. Abdullaev X. D. et al. EXPERIENCE OF USING THE DRUG GEPON IN THE TREATMENT OF PATIENTS FOCAL SCLERODERMA //ResearchJet Journal of Analysis and Inventions. – 2021. – Т. 2. – №. 03. – С. 60-63.

18. Абдуллаев Х. Д., Собиров М. С., Жумаева Д. Х. НЕРВНО-ПСИХИЧЕСКИЙ СТАТУС У БОЛЬНЫХ СЕБОРЕЕЙ // Молодежь и медицинская наука в XXI веке. – 2018. – С. 115-116.
19. Ахмедов Ш. К. и др. ЭФФЕКТИВНОСТЬ ЛЕЧЕНИЯ ИЗОТРЕТИНОИНА ПРИ ЛЕЧЕНИИ УГРЕВОЙ БОЛЕЗНИ // Академический журнал Западной Сибири. – 2015. – Т. 11. – №. 1. – С. 56-56.
20. Абдуллаев Х., Толибов М. Allergodermatozlar bilan bog'liq bo'lgan vulgar asneni kompleks davolash Samaraligini o'rganish // Журнал гепато-гастроэнтерологических исследований. – 2021. – Т. 2. – №. 3.2. – С. 73-74.
21. Davlatovich A. X., O'gli A. B. X., O'gli I. A. S. BOLALARDA GENITAL GERPESNI DAVOLASH SAMARADORLIGINI BAHOLASH // BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 367-369.
22. Абдуллаев Х. Д. и др. ЛАЗЕР В ЛЕЧЕНИИ ВИТИЛИГО // BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 495-500.
23. Xolmurodovich D. J., Umidovich N. T., Davlatovich A. X. CLINIC COURSE OF NONSPECIFIC PNEUMONIA // BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 510-513.
24. Davlatovich A. X. et al. USING IMMUNOMAX AND 0.1% TACROLIMUS OINTMENT IN THE TREATMENT OF VITILIGO // BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 559-562.
25. Davlatovich A. X. et al. EVALUATION OF THE GENERAL SOMATIC STATUS OF PATIENTS WITH VITILIGO BASED ON THE DETERMINATION OF THE CONTENT OF NATURAL ANTIBODIES TO VARIOUS ORGANIS AND TISSUES OF THE BODY // BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 472-476.
26. Davlatovich A. X. DETERMINATION OF GENE ACTIVITY IN VITILIGO PATIENTS // BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 451-454.
27. Xolmurodovich D. J., Orifovich R. S., Davlatovich A. X. FEATURES OF THE MICROELEMENT STATUS FOR THE DEVELOPMENT OF ATOPIC DERMATITIS IN CHILDREN // BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 447-450.
28. Hikmatovich I. N. et al. The use of Fungite in the Local Treatment of Genital Herpes // International Journal of Innovative Analyses and Emerging Technology. – 2021. – Т. 1. – №. 5. – С. 231-232.
29. Davlatovich A. X., Orifovich S. D., Baxtiyorovich A. S. Studying the Relationship of Local Immune Status and the Course of Burn Disease // Central Asian Journal of Medical and Natural Science. – 2022. – Т. 3. – №. 5. – С. 679-682.
30. Davlatovich A. X. VAGINAL TRIXOMONADLAR SHTAMMASINI TRIXOPOLGA VA XIMOTRIPSIN BILAN BILAN SEZGICHLIGINI ANIQLASH // BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 645-647.
31. Salamova L. A. et al. Evaluation of the effectiveness of the drug gepon in the treatment of genital candylomas. – 2021.